

QCD Field Strength Correlator at the One-Loop Order

V.I.Shevchenko*

*Institute for Theoretical and Experimental Physics
117218, B.Chermushkinskaya 25, Moscow, Russia*

Abstract

The leading perturbative contributions into the two-point gauge-invariant correlator $Tr\langle gF_{\mu\nu}(x)U(x,0)U(0,y)gF_{\rho\sigma}(y)U(y,0)U(0,x)\rangle$ are calculated at the one-loop order. It is shown, that nonlocal condensate $Tr\langle\alpha_s F_{\mu\nu}F_{\mu\nu}\rangle$ is nonzero at this order. The relation with the renormalization properties of Wilson loops is discussed.

*e-mail: shevchenko@vxitep.itep.ru

Our present theoretical understanding of strong interactions is based on quantum chromodynamics (QCD). It is well known, that the gauge nature of this theory leads to a wealth of its properties as well as to the great difficulties in the description of its dynamics. One source of these difficulties lies in the fact, that despite all observables one can calculate in the theory must be gauge invariant, the calculations actually involve noninvariant quantities like propagators at intermediate steps. The situation is studied well for the covariant gauges, at the same time for the noncovariant ones a lot of work still has to be done (see [1] and references therein). There is also a kind of temptation to consider one set of gauge conditions as "more physical" than another one supported by real simplifications arising then one chooses the "most adequate gauge" for the given problem (despite all physical observables are of course gauge independent).

Therefore it is very useful if possible to reformulate the theory in terms of gauge invariant quantities from the beginning. The latter are usually taken as Wilson loop functionals:

$$W(C) = \langle \text{Tr} \, P \exp \left(ig \int_C A_\mu dx^\mu \right) \rangle \quad (1)$$

The ordinary brackets denote usual average:

$$\langle O[A(x)] \rangle = \int DA(x) O[A(x)] \exp \left(-\frac{1}{4} \int d^4x F_{\mu\nu}^a F_{\mu\nu}^a \right)$$

where field strength tensor is $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$

Unfortunately, the loop approach [2] has two undesirable features. First of all, the original equations for loop variables are written for nonrenormalized quantities and complete renormalization program for them seems to be very difficult to proceed. Second, the loop equations are complicated since they defined not in the familiar coordinate space but in the loop space – subtle mathematical object itself.

All that makes attempts to look for an alternative set of gauge invariant quantities attractive. Recently a lot of work has been done (see [3] and references therein) in order to represent QCD on a way, where dynamical degrees of freedom are gauge invariant cumulants of the following type:

$$D_n = \langle \langle \prod_{i=1}^n G(x_i, x_0) \rangle \rangle \quad (2)$$

$$G(x_i, x_0) = U(x_0, x_i) g F_{\mu_i \nu_i}(x_i) U(x_i, x_0)$$

where nonabelian phase factors are

$$\begin{aligned} U(x, y) &= P \exp(-ig \int_x^y A_\mu(z) dz_\mu) = \\ &= 1 - ig \int_x^y A_\mu(z) dz_\mu + (-ig)^2 \int_x^y dt_\nu \int_x^t dz_\mu A_\mu(z) A_\nu(t) + \dots \end{aligned} \quad (3)$$

The double brackets in (2) denote the irreducible cumulants, for example $\langle \langle AB \rangle \rangle = \langle AB \rangle - \langle A \rangle \langle B \rangle$, see [4] and [8] for details.

Applying the nonabelian Stokes theorem [5] and cluster expansion property [4] one obtains:

$$\begin{aligned} W(C) &= \langle \text{Tr} P \exp(ig \int_{C=\partial S} A_\mu dx^\mu) \rangle = \langle \text{Tr} P \exp(i \int_S G(x_i, x_0) d\sigma(x_i)) \rangle = \\ &= \text{Tr} P \exp(i \sum_{n=1}^{\infty} D_n d^n \sigma) \end{aligned} \quad (4)$$

where D_n are defined in (2). It may be proven, that the r.h.s. of (4) does not depend on the reference point x_0 as it must be, if all D_n are taken into account. Therefore there is a natural question, which n are essential in the expansion (4). One easy notes, that for small contours, where the coupling constant is small, highest cumulants which are by definition non-reducible Green's functions are small too. Lattice calculations and some theoretical considerations demonstrate, that the same conclusion is going to hold in the

nonperturbative regime [6] and only a few lowest cumulants are important. In particular, the lowest nontrivial bilocal cumulant measured on the lattice in [7] is believed to be dominant and it is actually the case in the so called Gaussian model of the gluodynamics vacuum [3, 8].

Therefore it would be interesting to understand the perturbative behaviour of the lowest gauge-invariant functions (2). First of all, one could compare the renormalization properties of (2) and (1). It will be seen below, that they are different, inspite of the similarities encoded in (4). Second, such results may be useful for the lattice calculations, where perturbative part of the cumulants is measured at the small distances. It should also be noticed, that gluon propagator in the radial gauge is expressed as an integral of the quantities like (5) [1]. So analysing perturbative expansion of (5) we get some information about the perturbative series in Fock-Schwinger (radial) gauge.

In the present letter we are going to present leading perturbative contributions to the simplest $n = 2$ cumulant at the one-loop level:

$$D_{\mu\nu\rho\sigma}(x, y) = Tr \langle gF_{\mu\nu}(x)U(x, 0)U(0, y)gF_{\rho\sigma}(y)U(y, 0)U(0, x) \rangle \quad (5)$$

The integration in any phase factor is along the straight line, connecting its end points.

The detailed full calculation of this quantity in the limiting case $y = 0$ has been done recently in [9] by two methods – direct calculation of the corresponding Feynman diagrams and by methods of heavy quark effective theory. If the reference point ($x_0 = 0$ in our case) is chosen at one of the correlator arguments (i.e. points where the operators F act are connected by the straight line), the quantity (5) depends on the only vector $z = (y - x_0) - (x - x_0)$ and two different tensor structures can be introduced:

$$\delta_{\mu\rho}\delta_{\nu\sigma} - \delta_{\nu\rho}\delta_{\mu\sigma} = \Delta_{\mu\nu\rho\sigma}^{(1)} \quad (6)$$

and

$$\Delta_{\mu\nu\rho\sigma}^{(1)} - \frac{d}{2} \left(\frac{z_\mu z_\rho}{z^2} \delta_{\nu\sigma} - \frac{z_\nu z_\rho}{z^2} \delta_{\mu\sigma} + \frac{z_\nu z_\sigma}{z^2} \delta_{\mu\rho} - \frac{z_\mu z_\sigma}{z^2} \delta_{\nu\rho} \right) = \Delta_{\mu\nu\rho\sigma}^{(2)} \quad (7)$$

Note, that

$$\Delta_{\mu\nu\rho\sigma}^{(2)} \delta_{\mu\rho} \delta_{\nu\sigma} = 0$$

therefore only the part proportional to $\Delta^{(1)}$ contributing to the condensate $\langle \alpha_s F_{\mu\nu} F_{\mu\nu} \rangle$. Two scalar coefficient functions in front of $\Delta^{(1)}$ and $\Delta^{(2)}$:

$$D_{\mu\nu\rho\sigma}(x-y) = D(z) \Delta_{\mu\nu\rho\sigma}^{(1)} + D_1(z) \Delta_{\mu\nu\rho\sigma}^{(2)} \quad (8)$$

were actually calculated in [9] and appeared to be nonzero both contrary to the tree level results where only $D_1(z)$ presents.

Generally tensor structure of (5) is more complicated and the results of [9] should be recovered by taking the points x, y and $x_0 = 0$ lying on a line, for example $x = -y = z/2$. So our results can be used as an independent check of [9].

Let us briefly remind the situation in QED. The abelian analog of (5) was investigated in [10]. Calculation there simplifies greatly since phase factors are cancelled. Representing renormalized photon propagator in Feynman gauge as (we use Euclidean metric throughout the paper):

$$\langle eA_\mu(x) eA_\nu(y) \rangle = \delta_{\mu\nu} \frac{1}{4\pi^2} \frac{d(z^2)}{z^2} \quad (9)$$

where $z = y - x$ and $d(z^2) = e^2(z^2)$ - the renormalized effective charge and differentiating (9) with respect to z one easily finds:

$$\langle eF_{\mu\nu}(x) eF_{\rho\sigma}(y) \rangle = \frac{1}{\pi^2} \left[\Delta_{\mu\nu\rho\sigma}^{(2)} \left(\frac{d}{z^4} - \frac{d'}{z^2} + \frac{d''}{2} \right) - \Delta_{\mu\nu\rho\sigma}^{(1)} \frac{d''}{2} \right] \quad (10)$$

Here z should be taken nonzero in order to avoid contact terms and the derivative $d' \equiv de^2(z^2)/dz^2$. Only the term proportional to d'' contributes to $\langle F_{\mu\nu} F_{\mu\nu} \rangle$. Using the definition from [10] for the coordinate β -function :

$$z^2 \frac{de^2}{dz^2} = \beta(e^2)$$

it is easy to see that

$$\langle e^2 F_{\mu\nu}(x) F_{\mu\nu}(y) \rangle = \frac{6}{\pi^2} \frac{1}{z^4} \left(1 - \frac{d\beta(e^2)}{de^2} \right) \cdot \beta(e^2) \quad (11)$$

Taking $e^2(z^2) = e^2(\mu^2) + be^4(\mu^2) \ln[1/(\mu z)^2] + O(e^6)$ we find at the lowest nontrivial order:

$$\langle e^2 F_{\mu\nu}(x) F_{\mu\nu}(y) \rangle = -\frac{6be^4(\mu^2)}{\pi^2 z^4} \quad (12)$$

At the tree level in the $SU(N)$ Yang-Mills theory the only difference from abelian case in the common colour factor $(N^2 - 1)/2$:

$$D_{\mu\nu\rho\sigma}^{(0)} = \frac{N^2 - 1}{2} \frac{g^2}{\pi^2} \frac{1}{z^4} \cdot \Delta_{\mu\nu\rho\sigma}^{(2)} \quad (13)$$

Situation at the loop level is much more complicated in the nonabelian case. The main reason is the presence of phase factors $U(x, y)$ in the expression (5). Integrations over the straight lines, connecting the points x, y and $x_0 = 0$ lead to a new type of divergencies, additional to the usual ultraviolet ones. These divergencies are studied well in context of renormalization of the Wilson loops and we will explain below the role they play in renormalization of field correlators.

Let us proceed with the direct calculation of the leading g^4/ϵ terms in (5). The Feynman diagrams, contributing to the quantity under consideration are presented in Fig.1 and Fig.2. Naively only diagrams 1a, b, d and 2a, b, c, d, f, h give the leading contribution. Tree level contribution $D_{\mu\nu\rho\sigma}^{(0)}$ is defined in (13). Using dimensional regularization $d = 4 - 2\epsilon$ and keeping only $1/\epsilon$ terms we get in the Feynman gauge (we consider gluodynamics but one can include dynamical quarks easily):

$$\begin{aligned} 1a) + 1b) &= \frac{\alpha_s}{\pi} N \frac{5}{12} \left(\frac{1}{\epsilon} \right) \cdot D_{\mu\nu\rho\sigma}^{(0)} \\ 1d) &= -\frac{\alpha_s}{\pi} N \frac{3}{4} \left(\frac{1}{\epsilon} \right) \cdot D_{\mu\nu\rho\sigma}^{(0)} \end{aligned}$$

$$2b) + 2h) = \frac{\alpha_s}{\pi} N \left(\frac{1}{\epsilon} \right) \cdot D_{\mu\nu\rho\sigma}^{(0)}$$

$$2d) = \frac{\alpha_s}{\pi} N \left(\frac{1}{\epsilon} \right) \cdot \frac{1}{2} [(\pi - \gamma) \text{ctg}(\gamma)] \cdot D_{\mu\nu\rho\sigma}^{(0)}$$

In the last expression γ is an angle between x and y . It may be seen that diagram $2a$ does not contain singular parts and diagram $2f$ equals to zero. The straightforward check shows the coincidence with the results for the leading logarithmic terms from [9]. Note, that contrary to [9] we are working in the fundamental representation and our D from (5) is in fact $\mathcal{D}/2$ from [9].

All the above diagrams proportional to $\Delta^{(2)}$ and therefore do not contribute into the condensate

$$D_{\mu\nu\rho\sigma}(x, y) \cdot \delta_{\mu\rho} \delta_{\nu\sigma}$$

it is fulfilled by the only $2c$, the result is:

$$D_{\mu\nu\rho\sigma}(x, y) \cdot \delta_{\mu\rho} \delta_{\nu\sigma} =$$

$$= \frac{g^4}{16\pi^4} N(N^2 - 1) \left(\frac{1}{\epsilon} \right) \cdot \frac{1}{z^4} \left(\left[1 - 4 \frac{(xz)^2}{x^2 z^2} \right] + \left[1 - 4 \frac{(yz)^2}{y^2 z^2} \right] \right) \quad (14)$$

We note the nontrivial dependence on the angles between x , y and $z = y - x$, expressed via scalar products (xz) and (yz) . In particular case, if the special position of the points x , y and 0 is chosen the leading logarithmic contribution to the condensate is absent and only finite renormalization is needed. Namely, it is the case if the points x , y and 0 form the equal-side triangle of arbitrary size. This circumstance might be useful in the lattice calculations for fixing the normalization scale of the leading logarithms.

We see that the one-loop contribution to the field correlators cannot be entirely explained by charge renormalization. It is clearly indicated also in [9] where besides terms proportional to the first coefficient of the β -function there are additional terms in D_1 and D .

To proceed, let us compare the renormalization procedure for the Wilson loops (see [11] for details) and correlators. The former depends on whether the corresponding Wilson contour C is smooth or not despite the final results look in some sense similar. In the former case the average is renormalizable in the following sense:

$$W(C) = Z \cdot W_{ren}(C) \quad (15)$$

where infinite Z -factor contains only linear divergencies arising from the integrations over the contour:

$$Z \sim \exp \left(c \int_C dx_\mu \int_C dy_\nu D_{\mu\nu}(x-y) \right) \sim \exp \left(c \frac{L}{a} \right) \quad (16)$$

where L is the length of the contour a – ultraviolet cutoff and c – numerical constant not important for us here. At the same time logarithmic divergencies are absorbed in a usual way, so that renormalized Wilson average W_{ren} in the Eq. (15) is functional of the contour C and function of the renormalized coupling constant $g_{ren}(\mu)$ defined on the corresponding dynamical scale μ .

It is worth noting that a single diagram in the perturbative expansion of the r.h.s. of (15) typically gives contributions to both Z -factor and $W_{ren}(C)$, i.e. contain linearly and logarithmically divergent parts. Hence the factorization in the r.h.s. of (15) has double-faced nature - it is factorization of the diagrams together with the separation of contributions, coming from dangerous integration regions (i.e. linearly divergent terms) for each diagram itself.

As a result contour divergencies can be separated as a common Z -factor, which physically renormalizes the bare mass of the test particle, moving along the Wilson contour. This additional infinite constant in the exponent reflects the impossibility to observe the Wilson loop itself. At the same time if one extracts the physical quantities from the Wilson loop average then only logarithmic renormalization cause observable effect.

If the contour possesses cusps (and selfintersections) situation becomes more complicated [11]. Namely, each cusp leads to its own local Z – factor, depending on the cusp angle (but not on another characteristics of the contour). The Eq.(15) for the loop with K cusps having angles γ_i is to be modified in the following way:

$$W(C) = \prod_{i=1}^K Z_i(\gamma_i) \cdot W_{ren}(C) \quad (17)$$

The mixing between linear and logarithmic divergencies, mentioned above has some specific features for the loops with cusps. Namely, each cusp in addition to the linearly divergent factors, removed by the test particle mass renormalization introduces logarithmically divergent terms which have nothing to do with the coupling constant renormalization and must be subtracted by their own $Z(\gamma_i)$ –factors. There are two types of terms of this kind, which do depend on the cusp angle and which do not [11]. The latter plays important role in the perturbative expansion of the correlators. To see this, note, that the correlators can be easily obtained from the Wilson loop functional. Indeed, differentiating (1) with respect to $\delta\sigma_{\mu\nu}(x_i)$ and assuming that the contour C connects the points u, v with the reference point x_0 along the straight lines, one finds for the bilocal correlator:

$$\frac{1}{N_c} Tr g^2 \langle \langle F_{\mu\nu}(u, x_0) F_{\rho\sigma}(v, x_0) \rangle \rangle = - \frac{\delta^2 W(\tilde{C})}{\delta\sigma_{\mu\nu}(u) \delta\sigma_{\rho\sigma}(v)} \quad (18)$$

and analogous expressions for higher cumulants. Note, that area of the minimal surface bound by the contour $\tilde{C}$ from (18) is equal to zero and this contour generally has at least two cusps (typically even four, if the reference point x_0 does not coincide with one of the correlator arguments).

Comparing Eqs. (17) and (18) one concludes, that the derivatives in the r.h.s. of (18) must act on renormalized part of the Wilson loop as well as on the Z – factors. As a result mixing between different terms appears and

contour divergencies cannot be separated in some simple way as they can in the case of loops, so the perturbative behaviour of the correlators is controlled by both ultraviolet and contour logarithmic divergencies.

Let us do the final remark. It may be easily shown, that taking into account only the bilocal correlator in the cumulant expansion (4) one has perturbative contributions to the string tension at the one loop level since the kroneker part of (8) does not equal to zero. But area law is perturbatively impossible. The problem is cured by the contributions from the higher cumulants, in particular triple correlator $\langle FFF \rangle$ also contain such kroneker parts being considered at the same (g^4 in our case) order of perturbation theory [12]. As a result string tension is equal to zero in the field-strength formulation as it must be at any given order of perturbation theory if all cumulants contributing to string tension at this order are taken into account. Note in this respect that there are no terms proportional to the β -function coefficients in the function D [9] - it is a direct consequence of the fact that nonzero D has no physical sense in perturbation theory and all contributions into it must be cancelled by the next-order terms.

Acknowledgements

The work was supported by RFFI grants 96-02-19184 and RFFI-DFG 96-02-00088G. The author is grateful to the theory group of Heidelberg University where the part of the work was done for kind hospitality. Discussions with H.G.Dosch, M.Eidemüller, M.Jamin are gratefully acknowledged. The author would also like to thank Yu.A.Simonov for useful discussions and critical remarks and A.Vairo for pointing authors attention to the Ref.[6].

References

- [1] S. Leupold, H. Weigert, hep-th/9604015.

- [2] Yu.M. Makeenko, A.A. Migdal, Nucl.Phys. **B188** (1981) 269, for a review see A.A. Migdal, Phys.Rep. **102** (1983) 199.
- [3] H.G. Dosch, Prog.Part.Nucl.Phys. **33** (1994) 121.
H.G. Dosch, A. DiGiacomo and Yu.A. Simonov, in preparation.
- [4] N.G. Van Kampen, Stochastic Processes in Physics and Chemistry, North-Holland Physics Publishing, 1984.
- [5] Y. Aref'eva, Theor.Math.Phys. **43** (1980) 353.
- [6] G. Bali, N.Brambilla, A. Vairo, hep-lat/9709079, to appear in Phys.Lett.**B**.
L. Del Debbio, A. DiGiacomo, Yu.A. Simonov, Phys.Lett. **B 332** (1994) 111.
- [7] A. Di Giacomo and H. Panagopoulos, Phys. Lett. B 285 (1992) 133;
A. Di Giacomo, E.Meggiolaro and H. Panagopoulos, Nucl.Phys. **B483** (1997) 371.
- [8] H.G. Dosch, Phys. Lett. B 190 (1987) 177;
H.G. Dosch and Yu.A. Simonov, Phys. Lett. **B 205** (1988) 339;
- [9] M. Eidemüller , M. Jamin, hep-ph/9709419.
- [10] A.I. Vainshtein, V.I. Zakharov, Phys.Lett. **B225** (1989) 415.
- [11] V.S. Dotsenko, S.N. Vergeles, Nucl.Phys. **B169** (1980) 527;
R.A. Brandt, F. Neri, M.-A. Sato, Phys.Rev. **D24** (1981) 879.
- [12] V.I. Shevchenko, Yu.A. Simonov, in preparation.




Fig.1.




Fig.2.